

NUMERICAL DECOMPOSITION OF AFFINE ALGEBRAIC VARIETIES

SHAWKI AL-RASHED GERHARD PFISTER
RASHED@MATHEMATIK.UNI-KL.DE PFISTER@MATHEMATIK.UNI-KL.DE

DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF KAISERSLAUTERN,
ERWIN-SCHRÖDINGER-STR.,
67663 KAISERSLAUTERN,
GERMANY

ABSTRACT. An irreducible algebraic decomposition $\cup_{i=0}^d X_i = \cup_{i=0}^d (\cup_{j=1}^{d_i} X_{ij})$ of an affine algebraic variety X can be represented as an union of finite disjoint sets $\cup_{i=0}^d W_i = \cup_{i=0}^d (\cup_{j=1}^{d_i} W_{ij})$ called numerical irreducible decomposition (cf. [14],[15],[17],[18],[19],[21],[22],[23]). W_i corresponds to a pure i -dimensional X_i , and W_{ij} presents an i -dimensional irreducible component X_{ij} . Modifying this concepts by using partially Gröbner bases, local dimension, and the "Zero Sum Relation" we present in this paper an implementation in SINGULAR to compute the numerical irreducible decomposition. We will give some examples and timings, which show that the modified algorithms are more efficient if the number of variables is not too large. For a large number of variables BERTINI is more efficient. Note that each step of the numerical decomposition is parallelizable. For our comparisons we did not use the parallel version of BERTINI.

keyword: Witness point set, Homotopy function, Gröbner basis, Local dimension, Monodromy action, Zero Sum Relation.

1. INTRODUCTION

Given a system of n polynomials in $\mathbb{C}^N$,

$$f(x_1, \dots, x_N) := \begin{pmatrix} f_1(x_1, \dots, x_N) \\ \vdots \\ f_n(x_1, \dots, x_N) \end{pmatrix}.$$

Let $X=V(f)$ be the algebraic variety defined by the system above. X has an unique algebraic decomposition into d pure i -dimensional components X_i , $X = \cup_{i=0}^d X_i$. Where $X_i = \cup_{j=1}^{d_i} X_{ij}$ is the union of d_i i -dimensional irreducible components, $d_0, d_1, \dots, d_d$ positive integers.

The numerical irreducible decomposition (cf. [15],[17],[18],[19],[22]) is given as the union $W = \cup_{i=0}^d W_i = \cup_{i=0}^d (\cup_{j=1}^{d_i} W_{ij})$. The W_i are called i -Witness point sets and are given as an intersection of the pure i -dimensional component X_i of X with a generic linear space L in $\mathbb{C}^N$ of dimension $N-i$, the W_{ij} are called the irreducible

witness point sets presenting the irreducible components X_{ij} of dimension i with the following properties:

- W_{ij} consists of a finite number of points contained in X_{ij} .
- $\#(W_{ij}) = \deg(X_{ij})$ for $i \neq 0$.
- $W_{ij} \cap W_{il} = \emptyset$ for $j \neq l$.

Computing of the numerical irreducible decomposition uses numerical polynomial homotopy continuation methods(cf. [24],[25]). This requires that the number n of a given polynomial system has to be equal to the number N of the variables. Therefore we reduce the polynomial system which defines X to a square system of N polynomials in N variables (cf. [15],[17],[22]). Numerical irreducible decomposition (cf. [15],[17],[18],[19],[22]) is proceeded in three steps:

- 1st step reduces the polynomial system to a system of N polynomials in N variables and computes a finite set $\widehat{W}_i$ called witness point super set for a pure i -dimensional component X_i for $i=0,\dots,N-1$. $\widehat{W}_i$ consists of points on X_i and J_i a set of points on components of larger dimension the so-called Junk point set (cf. [15],[22]).
- 2nd step removes the points of J_i from $\widehat{W}_i$ to obtain a subset W_i of the pure i -dimensional component X_i (cf. [22]).
- 3rd step breakup W_i into irreducible witness point sets representing the i -dimensional irreducible components of X using two algorithms. The first algorithm finds points on the same irreducible component in the Witness point set connected by path tracking techniques applying the idea of monodromy. The second algorithm computes a linear trace for each component, which certifies the decomposition (cf. [14],[21]).

In the second section we present a modified algorithm to compute $\widehat{W}_i$ using the cascade algorithm (cf. [15],[17],[22]) and Gröbner bases in the zero-dimensional case. Then we use the homotopy function (cf. [22]), local dimension and Gröbner bases in the zero-dimensional case to remove Junk points from $\widehat{W}_i$ to obtain the i -Witness point set W_i . This is explained in third section. In the fourth section we explain how to use the "Zero Sum Relation" and monodromy action on the algebraic variety to breakup W_i into irreducible witness point sets. In the fifth section some examples are tested and timings are given on the basis of our SINGULAR implementation.

2. WITNESS POINT SUPER SET

Definition 2.1. Let $Z=V(f)$ be an affine algebraic variety in $\mathbb{C}^N$ of dimension d , and X be a pure i -dimensional component of Z . Let L_i be a generic linear space in $\mathbb{C}^N$ of dimension $N-i$. A set $\widehat{W}_i \subset \mathbb{C}^N$ is called i -Witness Point Super Set for X if it has the following properties:

- $\widehat{W}_i$ is a finite set of points.
- $X \cap L_i \subset \widehat{W}_i \subset Z \cap L_i$.

The union $\widehat{W}$ of all i -Witness Point Super Sets is called a Witness Point Super Set for Z .

In (cf. [15],[17],[22]) the cascade algorithm is used to compute $\widehat{W}_i$. It starts with $i=N-1$ to compute the Witness point Super Sets $\widehat{W}_i$. It needs to define a start

system $G(x)=0$ for the homotopy continuation method (cf. [24],[25]) and needs to know its solutions. We use a Gröbner basis to compute the dimension d of Z , then use the cascade algorithm (cf. [15],[17],[22]) which starts with $i=d-1$. We show below that we do not need to define a start system.

Algorithm 1 WITNESSPOINTSUPERSET

Input: $F_1, \dots, F_n \in \mathbb{C}[x_1, \dots, x_N]$.

Output: $\{f_1, \dots, f_N\}, \{\widehat{W}_r, \dots, \widehat{W}_d\}, L$. $\{f_1, \dots, f_N\}$ square system, $\widehat{W}_i$ Witness point super sets corresponding to a pure i -dimensional component of $V(f_1, \dots, f_N)$, L a set of linear polynomials defining a linear space of dimension $N-d$.

$f = \{f_1, \dots, f_N\}$ reduction of $F = \{F_1, \dots, F_n\}$ to a square system (cf. [15],[17],[22]);

$d = \dim(V(f_1, \dots, f_N))$, using Gröbner basis (cf. [8],[11]);

$r = N - \text{rank}(f)^1$, $\text{rank}(f)$ the rank of the Jacobian matrix of the system f at a generic point;

$L = \{l_1, \dots, l_d\}$ a set of d generic linear polynomials;

if $d = r$ **then**

 compute $T_d = V(f_1, \dots, f_N, l_1, \dots, l_d)$, using triangular sets, (cf. [8],[11]);

 set $\widehat{W}_d = \{(x_1, \dots, x_N) \mid (x_1, \dots, x_N) \in T_d, (x_1, \dots, x_N) \in V(F)\}$;

return $\{f_1, \dots, f_N\}, \{\widehat{W}_d\}, L$;

else

for $i = r$ **to** d **do**

if $i = 0$ **then**

$\Omega_i(f)(x) = f$;

else

$$\Omega_i(f)(x, z_1, \dots, z_i) =: \begin{pmatrix} f_1(x) + \sum_{j=1}^i \lambda_{1j} z_j \\ \vdots \\ f_N(x) + \sum_{j=1}^i \lambda_{Nj} z_j \\ l_1 + z_1 \\ \vdots \\ l_i + z_i \end{pmatrix}$$

$\lambda_{kj} \in \mathbb{C}$ generic, $k = 1, \dots, N, j = 1, \dots, i$;

for $i = d$ **to** r **do**

if $i = d$ **then**

 compute $T_i = V(\Omega_i(f)(x, z_1, \dots, z_i))$, using triangular sets, (cf. [8],[11]);

 set $\widehat{W}_i = \{(x_1, \dots, x_N) \mid (x_1, \dots, x_N, 0, \dots, 0) \in T_i, (x_1, \dots, x_N) \in V(F)\}$;

$S_i = T_i \setminus \{(x_1, \dots, x_N, z_1, \dots, z_i) \in T_i \mid z_1 = \dots = z_i = 0\}$;

else

 compute $T_i = V(\Omega_i(f)(x, z_1, \dots, z_i))$, using homotopy function with $\Omega_{i+1}(f)(x, z_1, \dots, z_i)$ as start system and S_{i+1} as start solution set (cf. [15],[17],[22]);

return $\{f_1, \dots, f_N\}, \{\widehat{W}_r, \dots, \widehat{W}_d\}, L$;

¹ $V(f_1, \dots, f_N)$ has no components of dimension smaller than $N - \text{rank}(f)$ (cf. [22]).

3. COMPUTATION WITNESS POINT SET

The witness point super set $\widehat{W}_i$ is an union of an i-Witness point set W_i and a junk point set J_i (cf. [15],[17],[22]),

$$\widehat{W}_i = W_i \cup J_i, \text{ where } W_i \subset X_i \text{ and } J_i \subset \cup_{j>i} X_j \text{ for } i = 0, 1, \dots, d.$$

We use Gröbner bases for the 0-dimensional ideal, local dimension and homotopy continuation method to remove the points of J_i from $\widehat{W}_i$ as follows:

Algorithm 2 WITNESSPOINTSET

Input: $\{f_1, \dots, f_N\} \subset \mathbb{C}[x_1, \dots, x_N]$, $\{\widehat{W}_r, \dots, \widehat{W}_d\}$ a list of witness point superset sets, $L = \{l_1, \dots, l_d\}$ a set of generic linear polynomials (Output of **Algorithm 1**).

Output: $\{f_1, \dots, f_N\}$, $\{W_r, \dots, W_d\}$, $L = \{l_1, \dots, l_d\}$. W_i a Witness point set corresponding to a pure i-dimensional component of $V(f_1, \dots, f_N)$.

$W_d = \widehat{W}_d$, $s_d = \#W_d$;

for $i = d - 1$ **to** r **do**

$W_i = \widehat{W}_i$;

for each point $w \in W_i$ **do**

 compute² $t = \dim_w Z$ for $Z = V(f_1 - f_1(w), \dots, f_N - f_N(w))$, using Gröbner basis (cf. [8],[11]);

if $t > i$ **then**

$W_i = W_i \setminus \{w\}$;

for each point $w \in W_i$ **do**

if $i = 0$ **then**

 choose $A \subset \mathbb{C}^{d \times N}$ a generic matrix and a generic $\epsilon \in \mathbb{C}^N$, $\|\epsilon\|$ small;

 compute $S = V(\{f_1, \dots, f_N, A(x - w)\})$, $T = V(\{f_1, \dots, f_N, A(x - w - \epsilon)\})$, using triangular sets (cf. [8],[11]);

if $\#S = \#T$ **then**

$W_i = W_i \setminus \{w\}$;

else

for $j = i + 1$ **to** d **do**

 choose $A \subset \mathbb{C}^{j \times N}$ a generic matrix;

if $j = d$ **then**

 compute $S = V(\{f_1, \dots, f_N, A(x - w)\})$, using triangular sets, (cf. [8],[11]);

if $\#S = s_d$ **then**

$W_i = W_i \setminus \{w\}$;

else

 compute $S = V(\{f_1, \dots, f_N, A(x - w)\})$, using homotopy function with start system $\{f_1, \dots, f_N, l_1, \dots, l_j\}$ and start solution W_j (cf. [22]);

if $w \in S$ **then**

$W_i = W_i \setminus \{w\}$;

return $\{f_1, \dots, f_N\}, \{W_r, \dots, W_d\}, L$;

² $t = \dim_w Z$ is less or equal to the local dimension of a point $v \in V(f_1, \dots, f_N)$ (cf. [9]) with $\|w - v\|$ small.

4. PARTITION WITNESS POINT SETS

In this section we show that the monodromy action on an algebraic variety Z and the Zero Sum Relation are sufficient to find the breakup of the k -Witness point set W_k into irreducible k -Witness point sets. We present here a modified version of the algorithms described in (cf. [14],[21]).

Let Z be a pure k -dimensional algebraic variety in $\mathbb{C}^N$, and $Z = \cup_{i=1}^r Z_i$ be the irreducible decomposition of Z . Let $\pi : \mathbb{C}^N \longrightarrow \mathbb{C}^k$ be a generic projection and let $l \subset \mathbb{C}^k$ be a general line.

Set


- $W_l := \pi^{-1}(l) \cap Z$ a set of r different curves in $\mathbb{C}^N$.
- U is a non-empty open subset of l consisting of all points $x \in l$ with $\pi^{-1}(x)$ transverse to Z .
- $W := \pi^{-1}(x) \cap Z$ for a generic element $x \in U$, and $V_x := V$ a subset of W .
- $W_i := \pi^{-1}(x) \cap Z_i$ for an irreducible k -dimensional component Z_i of Z .
- $\lambda : \mathbb{C}^N \longrightarrow \mathbb{C}$ a linear function one-to-one on W .

Define the function $s : U \longrightarrow \mathbb{C}$ by

$$s(y) = \sum_{z \in V_y} \lambda(z).$$

Where V_y is a subset of $\pi^{-1}(y) \cap Z$ defined by.

$$V_y := \{z \mid z \text{ on a curve through a point of } V\}.$$



Theorem 4.1. *Let l , U , W , W_i for $i=1, \dots, r$, and the functions λ , s be as above. If the function s is continuous and $V \cap W_i \neq \emptyset$ for some $i \in \{1, \dots, r\}$, then $W_i \subseteq V$.*

Example 4.1. *Before proving the theorem we illustrate it by an example.*

- Let Z be the algebraic variety of dimension one in $\mathbb{C}^2$ defined by the polynomial $f(x, y) = (x^2 + y^2 - 5)(x - 2y - 3)$. Let L_1 be the linear space of dimension one in $\mathbb{C}^2$ defined by the polynomial $l_1 = x + y - 3$.
- Define a homotopy function :

$$h(t, x(t), y(t)) := \begin{pmatrix} \alpha(t) \\ f(x(t), y(t)) \end{pmatrix}.$$

$\alpha : [0, 1] \longrightarrow p^{-1}(L)$ given by

$$\alpha(t) = (1 - t)l_0 + tl_1 = x + y - 2t - 1.$$

Let L_0 be the 1-dimensional linear space defined by the polynomial $l_0 = x + y - 1$, $L_0 \cap Z \neq \emptyset$. Let $G(N - k, N)$ be the Grassmanian and $R :=$

$\{(L_{N-k}, x) \in G(N-k, N) \times \overline{Z} \mid x \in L_{N-k} \cap \overline{Z}\}$ be the family of the intersections $L_{N-k} \cap \overline{Z}$, $L_{N-k} \subset \mathbb{P}^N$ k -dimensional linear spaces and $\overline{Z} \subset \mathbb{P}^N$ the closure of Z . Let $p : R \rightarrow G(N-k, N)$ be the canonical projection.

Then with conditions above $\alpha(t)$ maps a point in $L_1 \cap Z$ to a point in $L_0 \cap Z$ as t goes from 1 to 0.

Proof. (of theorem 4.1) Assume that $W_i \not\subseteq V$. Since $W_i \cap V \neq \emptyset$, then there are $a, b \in W_i$ such that a is not in V and $b \in V$. Let $a_1, \dots, a_r$ denote the points of the set $V \setminus \{b\}$. By (Corollary 3.5 in [14]) there is a loop α in the fundamental group $\pi_1(U, \pi^{-1}(x))$ with $\alpha(0) = \alpha(1)$ which takes a_j to a_j for all $j=1, \dots, r$, and interchanges a and b .

Since α is a continuous loop and $s : U \rightarrow \mathbb{C}$ is continuous, the composition $s \circ \alpha : [0, 1] \rightarrow \mathbb{C}$ is continuous and

$$s(\alpha(1)) = s(\alpha(0))$$

$$\lambda(a) + \sum_{j=1}^r \lambda(a_j) = \lambda(b) + \sum_{j=1}^r \lambda(a_j),$$

as t goes from 1 to 0. This implies that $\lambda(a) = \lambda(b)$. But this contradicts the fact that λ is one-to-one on W . Thus $W_i \subseteq V$. $\square$

Example 4.2. Let Z be a pure 1-dimensional component in $\mathbb{C}^2$ defined by the polynomial $f(x, y) = (y-x)(y-2x)(y-3x)$, and $Z = Z_1 \cup Z_2 \cup Z_3$ be the irreducible decomposition. Let $\pi : \mathbb{C}^2 \rightarrow \mathbb{C}$ be the projection given by $\pi(x, y) = x$, and $\lambda : \mathbb{C}^2 \rightarrow \mathbb{C}$, $\lambda(x, y) = y$.

Note that the restriction of π to Z , π_Z is proper and generically three-to-one with degree 3 equal to the degree of Z . λ is one-to-one on the fiber $\pi^{-1}(y) = \{(x, x), (x, 2x), (x, 3x)\}$. Let L be the linear space in the Grassmannian $G(1, 2)$ defined by the linear polynomial $l(x, y) = x + y - 2$. L intersects Z in the finite set $W := \{(1, 1), (\frac{2}{3}, \frac{4}{3}), (\frac{1}{2}, \frac{3}{2})\}$.

Let $V := \{(1, 1), (\frac{2}{3}, \frac{4}{3})\} \subset W$. The function $\sum_{v \in V} \lambda(v)$ given by $\lambda(x, x) + \lambda(x, 2x) = x + 2x = 3x$ is continuous. Then by the theorem above for an irreducible 1-Witness point set $W_1 \subset V$ it contains $\{(1, 1)\}$.

Preparation of the algorithm(IrrWitnessPointSet):

Let $Z_k = \cup_{i=1}^r Z_{ki}$ be the union of the irreducible k -dimensional components of the algebraic variety Z and L_k be the generic linear space in $\mathbb{C}^N$ defined by k linear equations

$$l_j = c_{j0} + c_{j1}x_1 + \dots + c_{jN}x_N.$$

for $j=1, \dots, k$ and $i=0, 1, \dots, N$, $c_{ij} \in \mathbb{C}$.

We use the generic linear space L_k to define the generic projection

$\pi : \mathbb{C}^N \rightarrow \mathbb{C}^{k+1}$, $\pi(x_1, \dots, x_N) := (z_1, \dots, z_k, z_{k+1})$ as follows:

$$\begin{pmatrix} x_1 \\ \vdots \\ x_N \end{pmatrix} \mapsto \begin{pmatrix} z_1 \\ \vdots \\ z_k \\ z_{k+1} \end{pmatrix} := \begin{pmatrix} c_{11} & c_{12} & \cdot & \cdot & c_{1N} \\ c_{21} & c_{22} & \cdot & \cdot & c_{2N} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ c_{k1} & c_{k2} & \cdot & \cdot & c_{kN} \\ p_1 & p_2 & \cdot & \cdot & p_N \end{pmatrix} \cdot \begin{pmatrix} x_1 \\ \vdots \\ \vdots \\ \vdots \\ x_N \end{pmatrix},$$

$p_1, \dots, p_N \in \mathbb{C}$ randomly chosen.

Set $\lambda(x_1, \dots, x_N) := z_{k+1}$ and $l := z_k \subset \mathbb{C}^k$ as in the theorem above. Let $y := c_{k0}$ vary and fix the other $c_{10}, \dots, c_{(k-1)0}$.

- Let $L_{k,y}$ be the linear spaces defined by the linear equations $l_1, \dots, l_{k-1}$ above and $l_{k,y} := y + c_{k1}x_1 + \dots + c_{kN}x_N$.
- For the subset V_y of $W_y := L_{K,y} \cap Z_k$ the function s is given as a function $s_y : Z_k \cap L_{k,y} \rightarrow \mathbb{C}$,

$$s_y(x_1, \dots, x_N) := \sum_{(x_1, \dots, x_N) \in V_y} \lambda(x_1, \dots, x_N).$$

It is convenient to test the linearity of s_y .

- Since $y = -(c_{k1}x_1 + \dots + c_{kN}x_N)$, then we can define a function s

$$s : \mathbb{C} \rightarrow \mathbb{C}, s(y) := s_y(x_1, \dots, x_N) = \sum_{(x_1, \dots, x_N) \in V_y} \lambda(x_1, \dots, x_N).$$

- s_y is a linear function in $x = (x_1, \dots, x_N)$ if and only if s is linear in y .
- To test the linearity of s , we take three values of y in $\mathbb{C}$, say a, b, c .

If there exist $A, B \in \mathbb{C}$ such that:

$$(s(a) = Aa + B, s(b) = Ab + B) \implies s(c) = Ac + B. \quad (4.1)$$

Then s is linear in y .

Here $s(a)$, $s(b)$ and $s(c)$ correspond to the subsets $V_a \subset W_a = Z_k \cap L_{k,a}$, $V_b \subset W_b = Z_k \cap L_{k,b}$ and $V_c \subset W_c = Z_k \cap L_{k,c}$ respectively with $\#V_a = \#V_b = \#V_c = m$.

So far this is the approach which can be found in [14]. Now we give some modifications.

- The condition (4.1) of the linearity above is equivalent to the following equation

$$s(a)(b - c) + s(b)(c - a) + s(c)(a - b) = 0. \quad (4.2)$$

- From **Theorem 4.1** we obtain: If $W_{kj} \cap V_a \neq \emptyset$ for some $j \in \{1, \dots, r\}$ and the condition (4.1) of the linearity above is true, then $W_{kj} \subseteq V_a$.
- Let $Z(y) := \{z = \sum_{t=1}^N p_t v_t \mid v = (v_1, \dots, v_N) \in V_y, p = (p_1, \dots, p_N) \in \mathbb{C}^N\}$. Then

$$s(y) = \sum_{v \in V_y} \lambda(v) = \sum_{v \in V_y} \left(\sum_{t=1}^N p_t v_t \right) = \sum_{z \in Z(y)} z.$$

- To compute the sets V_b and V_c we use the homotopy function as t goes from 1 to 0 using V_a as start set. In particular

$$V_b := ((1 - t)L_{k,b} + tL_{k,a}) \cap Z, V_c := ((1 - t)L_{k,c} + tL_{k,a}) \cap Z, \text{ as } t \text{ goes } 1 \rightarrow 0.$$

Continuation of the homotopy function implies that the i -th points in the sets V_a , V_b and V_c are on the same irreducible component.

- Let $V_a := \{v_1, \dots, v_m\}$, $V_b := \{\bar{v}_1, \dots, \bar{v}_m\}$ and $V_c := \{\hat{v}_1, \dots, \hat{v}_m\}$ be the sets computed by using the homotopy function above. Let $Z(a) := \{a_1, \dots, a_m\}$, $Z(b) := \{b_1, \dots, b_m\}$ and $Z(c) := \{c_1, \dots, c_m\}$ be the sets corresponding to the set V_a , V_b and V_c respectively and defined as $Z(y)$ above.

From (4.2) we get a condition equivalent to the condition (4.1) of the linearity

$$(b - c) \sum_{i=1}^m a_i + (c - a) \sum_{i=1}^m b_i + (a - b) \sum_{i=1}^m c_i = 0. \quad (4.3)$$

The condition (4.3) is called **Zero Sum Relation** (cf. [7]) of a given subset $V_a \subseteq W$ denoted by $ZSR(V_a)$.

- The sets V_a , V_b and V_c have distinct points and the same cardinality m , then obviously

$$ZSR(V_a) = \sum_{a_i \in V_a} ZSR(\{a_i\}). \quad (4.4)$$

where $ZSR(\{a_i\}) = (b - c)a_i + (c - a)b_i + (a - b)c_i$ is defined as a Zero Sum Relation of a given point in V_a .

Algorithm 3 IRRWITNESSPOINTSET

Input: $\{f_1, \dots, f_N\} \subset \mathbb{C}[x_1, \dots, x_N]$, $\{W_r, \dots, W_d\}$, a list of witness point sets, $L = \{l_1, \dots, l_d\}$ a set of generic linear polynomials (Output of **Algorithm 2**). Where $W_k = \{w_1, \dots, w_{m_k}\}$ are witness point sets for a pure k -dimensional component Z_k of $Z = V(f_1, \dots, f_N)$, $k = r, \dots, d$.

Output: $\{\{W_{r1}, \dots, W_{rt_r}\}, \dots, \{W_{d1}, \dots, W_{dt_d}\}\}$, W_{kr_k} irreducible Witness point sets corresponding to a k -dimensional irreducible component Z_{kr_k} of Z_k .

for $k = r$ **to** d **do**

$a := c_{k0}$, define L_{ka} to be the linear space defined by the subset $\{l_1, \dots, l_k\} \subset L$.

choose $b, c \in \mathbb{C}$ generic, define L_{kb}, L_{kc} as above;

$W_a = W_k$, $W_b = \emptyset$, $W_c = \emptyset$, $R = \emptyset$;

choose $p_1, \dots, p_N \in \mathbb{C}$;

for $i = 1$ **to** m_k **do**

compute $\{v_i\} \subset Z \cap L_{k,b}$ and $\{\widehat{v}_i\} \subset Z \cap L_{k,c}$ using the homotopy function with $\{f_1, \dots, f_N, l_1, \dots, l_{k-1}, l_{k,a}\}$ as start system and $\{w_i\}$ as start solution; compute the Zero Sum Relation of $\{w_i\}$:

$$r_i = (a - b) \left(\sum_{j=1}^N p_j \widehat{v}_{ij} \right) + (b - c) \left(\sum_{j=1}^N p_j w_{ij} \right) + (c - a) \left(\sum_{j=1}^N p_j v_{ij} \right);$$

$R = R \cup \{r_i\}$ ³;

int $t_k = 0$;

while $R \neq \emptyset$ **do**

if $\sum_{t \in T} t = 0$ and T is a smallest subset⁴ of R **then**

$t_k = t_k + 1$;

$W_{kt_k} \subset W_a$ consists of the points corresponding of the points of T ;

$R = R \setminus T$;

return $\{\{W_{r1}, \dots, W_{rt_r}\}, \dots, \{W_{d1}, \dots, W_{dt_d}\}\}$;

³the $i - th$ point in R corresponds to the $i - th$ point in W_a ;

⁴smallest subset with respect to the cardinality.

We give an example of a pure 2-dimensional variety Z which is an union of two 2-dimensional irreducible components Z_1 and Z_2 . Z_1 is of degree three and Z_2 is of degree two. The 2-Witness point set W for Z is given as a finite subset of Z consisting of five points $\{w_1, w_2, w_3, w_4, w_5\}$. Z_1 should contain three points $W_1 := \{w_1, w_2, w_3\}$ and the remaining points $W_2 := \{w_4, w_5\}$ are on Z_2 . The algorithms (cf. [14],[21]) use the homotopy function at least nine times to breakup W into W_1 and W_2 . We will show below that we do not need more than five times to use the homotopy function to breakup W into W_1 and W_2 .

Example 4.3.

Let Z be the algebraic variety of dimension two in $\mathbb{C}^3$ defined by the polynomial $f(x, y, z) = (x^3 + z)(x^2 - y)$. Let L be the linear space of dimension one in $\mathbb{C}^3$ defined by the linear equations $l_1 = 4x + 7y + 2z + 6$, $l_2 = 5x + 7y + 3z + 6$. Then $W := L \cap Z = \{w_1, w_2, w_3, w_4, w_5\}$, where⁵ $w_1 = (1, -1.1428571429, -1)$, $w_2 = (0, -0.8571428571, 0)$, $w_3 = (-0.1428571429 + i * 0.9147320339, -0.8163265306 - i * 0.2613520097, 0.1428571429 - i * 0.9147320339)$, $w_4 = (-1, -0.5714285714, 1)$, $w_5 = (-0.1428571429 - i * 0.9147320339, -0.8163265306 + i * 0.2613520097, 0.1428571429 + i * 0.9147320339)$.

We now illustrate *Algorithm3 (IrrWitnessPointSet)*:

- Use the linear space L_1 to define the linear projection $\pi : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ as follows

$$\pi(x, y, z) := \begin{pmatrix} 4 & 7 & 2 \\ 5 & 7 & 3 \\ 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = (4x + 7y + 2z, 5x + 7y + 3z, x + 2y + 3z).$$

- Define the linear space $L_{1,c}$ of dimension one in $\mathbb{C}^3$ by the linear equations $l_1 = 4x + 7y + 2z + 6$, $l_c = 5x + 7y + 3z + c$, where c is generically chosen in $\mathbb{C}$. Then

$$\pi_{Z \cap L_{1,c}}(x, y) = (-6, -c, x + 2y + 3z).$$

- Define the linear function $\lambda : \mathbb{C}^2 \rightarrow \mathbb{C}$ by $\lambda(x, y, z) := x + 2y + 3z$.
- For $a=6$, let $V_1 = V_a := \{w_{11} = (1, -1.1428571429, -1)\} \subset W$, $L_{1,a} := L$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_a = 5x + 7y + 3z + 6$. Then⁶ $Z(a) = \{\sum_{v \in V_a} \lambda(v) = w_{11}[1] + 2(w_{11}[2]) + 3(w_{11}[3])\} = \{-4.2857142858\}$.
- Let $b=9$, $L_{1,b}$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_b = 5x + 7y + 3z + 9$. Compute $V_b := (tL_{1,a} + (1-t)L_{1,b}) \cap Z = \{w_{12} = (1.671699881657157, -0.4776285376163331, -4.671699881657164)\}$ as t goes from 1 to 0, using V_a as a start solution. $Z(b) = \{w_{12}[1] + 2(w_{12}[2]) + 3(w_{12}[3])\} = \{-13.2986568385470012\}$.
- Let $c=63$, $L_{1,c}$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_c = 5x + 7y + 3z + 63$. Compute $V_c := (tL_{1,a} + (1-t)L_{1,c}) \cap Z = \{w_{13} = (3.935100643260828, 14.30425695906836, -60.93510064326094)\}$ as t goes

⁵Note that the values of w_i are approximate values. The following equalities are therefore to interpret as approximations of the points w_i .

⁶we use the notation $w_{ij} = (w_{ij}[1], w_{ij}[2], w_{ij}[3])$ for $i=1, \dots, 5, j=1, 2, 3$.

from 1 to 0, using V_a as a start solution. $Z(c) = \{w_{13}[1] + 2(w_{13}[2]) + 3(w_{13}[3])\} = \{-150.261687368385272\}$.

Zero Sum Relation of $V_1 = \{(1, -1.1428571429, -1)\}$:

$$\begin{aligned} r_1 &:= \sum_{a \in Z(a)} (b - c) + \sum_{b \in Z(b)} (c - a) + \sum_{c \in Z(c)} (a - b) = \\ &= -75.8098062588232524. \end{aligned}$$

The zero sum relation set of $V_1 = \{(1, -1.1428571429, -1)\}$ is $R_1 := \{r_1 = -75.8098062588232524\}$.

- Let $a=6$, $V_a := \{w_{11} = (0, -0.8571428571, 0)\} \subset W$, $L_{1,a} := L$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_a = 5x + 7y + 3z + 6$. Then $Z(a) = \{\sum_{v \in V_a} \lambda(v) = w_{11}[1] + 2(w_{11}[2]) + 3(w_{11}[3])\} = \{-1.7142857142\}$.
- Let $b=9$, $L_{1,b}$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_b = 5x + 7y + 3z + 9$. Compute $V_b := (tL_{1,a} + (1-t)L_{1,b}) \cap Z = \{w_{12} = (-0.8358499408285809 + i * 1.046869318849985, 0.2388142688081706 - i * 0.2991055196714253, -2.164150059171436 - i * 1.046869318849981)\}$ as t goes from 1 to 0, using V_a as a start solution. $Z(b) = \{w_{12}[1] + 2(w_{12}[2]) + 3(w_{12}[3])\} = \{-6.8506715807265477 - i * 2.6919496770428086\}$.
- Let $c=63$, $L_{1,c}$ the linear space defined by $l_1 = 4x + 7y + 2z + 6$, $l_c = 5x + 7y + 3z + 63$. Compute $V_c := (tL_{1,a} + (1-t)L_{1,c}) \cap Z = \{w_{13} = (-1.967550321630417 + i * 3.257877039491183, 15.99072866332302 - i * 0.9308220112831772, -55.03244967836969 - i * 3.257877039491242)\}$ as t goes from 1 to 0, using V_a as a start solution. $Z(c) = \{w_{13}[1] + 2(w_{13}[2]) + 3(w_{13}[3])\} = \{-135.083442030093447 - i * 8.3773981015488974\}$.

Zero Sum Relation of $V_2 = \{(0, -0.8571428571, 0)\}$:

$$\begin{aligned} r_2 &:= \sum_{a \in Z(a)} (b - c) + \sum_{b \in Z(b)} (c - a) + \sum_{c \in Z(c)} (a - b) = \\ &= 107.3334745556671221 - i * 128.308937286793398. \end{aligned}$$

The zero sum relation set of $V_2 = \{(0, -0.8571428571, 0)\}$ is

$R_2 := \{r_2 = 107.3334745556671221 - i * 128.308937286793398\}$.

- For the other points $V_3 = \{w_3\}$, $V_4 = \{w_4\}$ and $V_5 = \{w_5\}$, we found the zero sum relations $R_3 := \{r_3 = -9.38237104997583366 + i * 127.0170767088\}$, $R_4 := \{r_4 = -31.5236682999307779 + i * 128.3089372867945956\}$ and $R_5 := \{r_5 = 9.382371038077068 - i * 127.0170767088\}$.
- The set of Zero Sum Relation for all points of W is $R = \cup_{j=1}^5 R_j = \{r_1, r_2, r_3, r_4, r_5\}$, where i -th point in W corresponds i -th point in R .
- Find the smallest subset T of R with $\sum_{t \in T} t = 0$, which corresponds an irreducible Witness point set of W . Then we get $T_1 = \{r_3, r_5\}$, $T_2 = \{r_1, r_2, r_4\}$ corresponding to the irreducible Witness point sets $W_1 = \{w_3, w_5\}$, $W_2 = \{w_1, w_2, w_4\}$ respectively.

Remark 4.1. The points of a Witness point set are computed approximately by using the homotopy continuation method. Therefore the result of the Zero Sum Relation is only almost zero.

5. EXAMPLES AND TIMINGS WITH SINGULAR AND BERTINI

In this section we provide examples with timings of the algorithms `WitnessPointSuperSet`, `WitnessPointSet`, and `IrrWitnessPointSet` implemented in SINGULAR to compute the numerical decomposition of a given algebraic variety defined by a polynomial system and compare them with the results of BERTINI . We did not use the parallel features of BERTINI .

We tested to versions of the implementations in BERTINI using the cascade algorithm and using the regenerative cascade algorithm. Timings are conducted by using the 32-bit version of SINGULAR 3-1-1 (cf. [8]) and BERTINI 1.2 (cf. [3]) on an Intel® Core(TM)2 Duo CPU P8400 @ 2.26 GHz 2.27 GHz, 4 GB RAM under the Kubuntu Linux operating system.

Let Z be the algebraic variety defined by the following polynomial system:

Example 5.1. (cf. [17]).

$$f(x, y, z) = \begin{pmatrix} (y - x^2)(x^2 + y^2 + z^2 - 1)(x - \frac{1}{2}) \\ (z - x^3)(x^2 + y^2 + z^2 - 1)(y - \frac{1}{2}) \\ (y - x^2)(z - x^3)(x^2 + y^2 + z^2 - 1)(z - \frac{1}{2}) \end{pmatrix}$$

Example 5.2. (cf. [22], Example 13.6.4).

$$f(x, y, z) = \begin{pmatrix} x(y^2 - x^3)(x - 1) \\ x(y^2 - x^3)(y - 2)(3x + y) \end{pmatrix}$$

Example 5.3.

$$f(x, y, z) = \begin{pmatrix} (x^3 + z)(x^2 - y) \\ (x^3 + y)(x^2 - z) \\ (x^3 + z)(x^3 + y)(z^2 - y) \end{pmatrix}$$

Example 5.4.

$$f(x, y, z) = \begin{pmatrix} x(y^2 - x^3)(x - 1) \\ x(3x + y)(y^2 - x^3)(y - 2) \\ x(y^2 - x^3)(x^2 - y) \end{pmatrix}$$

Example 5.5.

$$f(x, y, z) = \begin{pmatrix} (x - 1)((x^3 + z) + (x^2 - y)) \\ (x^3 + z)(x^2 - y) \\ (x^3 + z)(x^2 - 1) \end{pmatrix}$$

Example 5.6.

$$f(x, y, z) = \begin{pmatrix} (y - x^2)(x^2 + y^2 + z^2 - 1)(x - \frac{1}{2}) + x^5 \\ (z - x^3)(x^2 + y^2 + z^2 - 1)(y - \frac{1}{2}) + y^4 \\ (y - x^2)(z - x^3)(x^2 + y^2 + z^2 - 1)(z - \frac{1}{2}) + z^6 \end{pmatrix}$$

Example 5.7.

$$f(x, y, z) = \begin{pmatrix} x(y^2 - x^3)(x - 1) + y^2 \\ x(y^2 - x^3)(y - 2)(3x + y) + x^3 \end{pmatrix}$$

Example 5.8.

$$f(x, y, z) = \begin{pmatrix} (x^3 + z)(x^2 - y) + x^4 \\ (x^3 + y)(x^2 - z) + y^3 \\ (x^3 + z)(x^3 + y)(z^2 - y) + z^5 \end{pmatrix}$$

Example 5.9.

$$f(x, y, z) = \begin{pmatrix} f_1 = -3568891411860300072x^5 + 1948764938x^4 + \\ 3568891411860300072x^2y^2 - 1948764938xy^2 \\ f_2 = -5105200242937540320x^5y - 1701733414312513440x^4y^2 + \\ 11692589628x^5 + 3897529876x^4y + 5105200242937540320x^2y^3 + \\ 1701733414312513440xy^4 - 11692589628x^2y^2 - 3897529876xy^3 \end{pmatrix}$$

Example 5.10.

$$f(x, y, z) = \begin{pmatrix} f_1 = -356737285367005125x^5 - 92300457164036000x^3y + \\ 1121648050080163317x^2z + 290209720279281056yz \\ f_2 = -356737285367005125x^5 + 887060318883271500x^3z + \\ 1121648050080163317x^2y - 2789081819567309964yz \\ f_3 = -356737285367005125x^5z^2 + 356737285367005125x^5y + \\ 887060318883271500x^3z^3 - 887060318883271500x^3yz + \\ 1121648050080163317x^2z^3 - 1121648050080163317x^2yz - \\ 2789081819567309964z^4 + 2789081819567309964yz^2 \end{pmatrix}$$

Example 5.11.

$$f(x, y, z) = \begin{pmatrix} f_1 = x^5y^2 + 2x^3y^4 + xy^6 + 2x^3y^2z^2 + 2xy^4z^2 + xy^2z^4 - x^4y^2 \\ - 2x^2y^4 - y^6 - x^5z - 2x^3y^2z - xy^4z - 2x^2y^2z^2 - 2y^4z^2 - \\ 2x^3z^3 - 2xy^2z^3 - y^2z^4 - xz^5 - 3x^3y^2 - 3xy^4 + x^4z + \\ 2x^2y^2z + y^4z - 3xy^2z^2 + 2x^2z^3 + 2y^2z^3 + z^5 + 3x^2y^2 + \\ 3y^4 + 3x^3z + 3xy^2z + 3y^2z^2 + 3xz^3 + 2xy^2 - 3x^2z - 3y^2z - \\ 3z^3 - 2y^2 - 2xz + 2z \\ f_2 = x^6y + 2x^4y^3 + x^2y^5 + 2x^4yz^2 + 2x^2y^3z^2 + x^2yz^4 - \\ 5x^6 - 10x^4y^2 - 5x^2y^4 - x^4yz - 2x^2y^3z - y^5z - 10x^4z^2 - \\ 10x^2y^2z^2 - 2x^2yz^3 - 2y^3z^3 - 5x^2z^4 - yz^5 - 3x^4y - \\ 3x^2y^3 + 5x^4z + 10x^2y^2z + 5y^4z - 3x^2yz^2 + 10x^2z^3 + \\ 10y^2z^3 + 5z^5 + 15x^4 + 15x^2y^2 + 3x^2yz + 3y^3z + 15x^2z^2 \\ + 3yz^3 + 2x^2y - 15x^2z - 15y^2z - 15z^3 - 10x^2 - 2yz + 10z \\ f_3 = x^6y^2z + 2x^4y^4z + x^2y^6z + 2x^4y^2z^3 + 2x^2y^4z^3 + \\ x^2y^2z^5 - 7x^6y^2 - 14x^4y^4 - 7x^2y^6 - x^6z^2 - 17x^4y^2z^2 - \\ 17x^2y^4z^2 - y^6z^2 - 2x^4z^4 - 11x^2y^2z^4 - 2y^4z^4 - x^2z^6 - \\ y^2z^6 + 7x^6z + 18x^4y^2z + 18x^2y^4z + 7y^6z + 15x^4z^3 + \\ 27x^2y^2z^3 + 15y^4z^3 + 9x^2z^5 + 9y^2z^5 + z^7 + 21x^4y^2 + \\ 21x^2y^4 - 4x^4z^2 + 13x^2y^2z^2 - 4y^4z^2 - 11x^2z^4 - 11y^2z^4 - \\ 7z^6 - 21x^4z - 40x^2y^2z - 21y^4z - 24x^2z^3 - 24y^2z^3 - 3z^5 - \\ 14x^2y^2 + 19x^2z^2 + 19y^2z^2 + 21z^4 + 14x^2z + 14y^2z + \\ 2z^3 - 14z^2 \end{pmatrix}$$

Example 5.12.

$$f(x_1, x_2, x_3, x_4, x_5) = \begin{pmatrix} x_5^2 + x_1 + x_2 + x_3 + x_4 - x_5 - 4 \\ x_4^2 + x_1 + x_2 + x_3 - x_4 + x_5 - 4 \\ x_3^2 + x_1 + x_2 - x_3 + x_4 + x_5 - 4 \\ x_2^2 + x_1 - x_2 + x_3 + x_4 + x_5 - 4 \\ x_1^2 - x_1 + x_2 + x_3 + x_4 + x_5 - 4 \end{pmatrix}$$

Example 5.13.

$$f(a, b, c, d, e, f, g) = \begin{pmatrix} a^2 + 2de + 2cf + 2bg + a \\ 2ab + e^2 + 2df + 2cg + b \\ b^2 + 2ac + 2ef + 2dg + c \\ 2bc + 2ad + f^2 + 2eg + d \\ c^2 + 2bd + 2ae + 2fg + e \\ 2cd + 2be + 2af + g^2 + f \\ d^2 + 2ce + 2bf + 2ag + g \end{pmatrix}$$

Example 5.14. . *cyclic 4-roots problem.*(cf.[5],[6]).

Example 5.15. . *cyclic 5-roots problem.*(cf.[5],[6]).

Example 5.16. . *cyclic 6-roots problem.*(cf.[5],[6]).

Example 5.17. . *cyclic 7-roots problem.*(cf.[5],[6]).

Example 5.18. . *cyclic 8-roots problem.*(cf.[5],[6]).

Example 5.19. .

$$f(x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{21}, x_{22}, x_{23}, x_{24}, x_{25}, x_{31}, x_{32}, x_{33}, x_{34}, x_{35}) =$$

$$= \begin{pmatrix} -x_{12}x_{21} + x_{11}x_{22} \\ -x_{13}x_{22} + x_{12}x_{23} \\ -x_{14}x_{23} + x_{13}x_{24} \\ -x_{15}x_{24} + x_{14}x_{25} \\ -x_{22}x_{31} + x_{21}x_{32} \\ -x_{23}x_{32} + x_{22}x_{33} \\ -x_{24}x_{33} + x_{23}x_{34} \\ -x_{25}x_{34} + x_{24}x_{35} \end{pmatrix}$$

Table 1 summarizes the results of the timings to compute the numerical decomposition⁷.

Remark 5.1. *The timings show that for an increasing number of variables the original method of (cf.[14],[15],[17],[21],[22]) becomes more efficient. One reason is that the computation of triangular sets which is used in SINGULAR for solving polynomial systems is expensive in this case. Therefore the **Algorithm1**, **Algorithm2** become slow in this situation. This is not true for **Algorithm3**.*

Replacing the solving of polynomial system using triangular sets by homotopy function methods but keeping the computation of the dimension and starting in this dimension is more efficient in a case of a large number of variables.

⁷(re) means using the regenerative cascade algorithm instead of the cascade algorithm

Example	BERTINI	BERTINI (re)	SINGULAR
5.1	134.45s	39s	36.07
5.2	3.08s	2.5s	1.49s
5.3	1min 21.28s	27.4s	4.02s
5.4	18.56s	2.7s	1.77s
5.5	15.36s	8.6s	1.29s
5.6	4min 13s	15min 2s	2min 27s
5.7	1.83s	1.6s	0.39s
5.8	3min 29s	10min 43s	1.69s
5.9	16s	7s	2s
5.10	2min 57s	28s	2min 35s
5.11	44min 56s	2min 37s	4min 3s
5.12	4.73s	6s	0.37s
5.13	5.84s	8s	1s
5.14	1.43s	4.3s	0.79s
5.15	3.54s	10s	0.57s
5.16	3min 23.26s	2min 29s	1.43s
5.17	2h 11min 57s	32min 17s	stopped after 5h
5.18	19h48min 17s	6h45min2s	stopped after 50h
5.19	1min 57s	51s	stopped after 3h

TABLE 1. Total running times for the computing a numerical decomposition of the examples above

REFERENCES

- [1] E. L. Allgower and K. George. *Numerical Continuation Methods, an Introduction*, Springer Series in Comput. Math, Vol. 13, Springer- Veralg, Berlin, Heidelberg, New Yourk, 1990.
- [2] E. Arbarello, M. Coranalba, P.A. Griffiths and J. Harris. *Geometry of Algebraic Curves*, volume I. Volume 267 of *Grundlehren Math. Wiss.*, Springer Verlag, New York, 1985.
- [3] D. J. Bates, J. D. Hauenstein, A. J. Sommese and C. W. Wampler . BERTINI : *Software for Numerical Algebraic Geometry*. <http://www.nd.edu/~sommese/bertini/>.
- [4] S. C. Billups, A. P. Morgan and L. T. Watson . Algorithm 652. Hompack: *A Suite codes for Globally Convergent Homotopy Algorithms*. ACM Transactions on Mathematical Software, Vol.13. No. 3, Septemper 1987, Pages 281- 310.
- [5] G. Björck and R. Fröberg. *A faster way to count the solutions of inhomogeneous systems of algebraic equations, with applications to cyclic n-roots*. *J. Symbolic Computation*, 12:329–336, 1991.
- [6] G. Björck. *Functions of modulus one on Z_n whose Fourier transforms have constant modulus, and cyclic n-roots*. In: J.S. Byrnes and J.F. Byrnes, Editors, *Recent Advances in Fourier Analysis and its Applications* 315, NATO Adv. Sci. Inst. Ser. C. Math. Phys. Sci., Kluwer (1989), pp. 131-140.
- [7] R. M. Corless, A. Galligo, I. S. Kotsireas and S. M. Watt. *A Geometric- Numeric Algorithm for Absolute Factorization of Multivariate Polynomials*. ISSAC 2002, July 7- 10, 2002, Lille, France.
- [8] W. Decker, G.-M Greuel, G. Pfister and H. Schönemann. SINGULAR 3-1-1 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2010).
- [9] G. Fischer. *Complex Alalytic Geometry*, Lecture Notes in Mathematics 538 (1976).
- [10] W. Fulton. *Intersection Theory*, volume (3) 2 of *Ergeb. Math. Grenzgeb*. Springer Verlag, Berlin, 1984.
- [11] G.-M. Greuel and G.Pfister. *A SINGULAR Introduction to Commutative Algebra*. Second edition, Springer (2007).

- [12] A. P. Morgan and L. T. Watson. *A globally convergent parallel algorithm for zeros of polynomial systems*. Nonlinear Anal. (1989), 13(11), 1339–1350.
- [13] A. P. Morgan and A.J. Sommese. *Computing all solutions to polynomial systems using homotopy continuation*. Appl. Math. Comput., 115- 138. Errata: Appl. Math. Comput. 51 (1992), p. 209. Nonlinear Anal.
- [14] A.J. Sommese, J Verschelde and C.W. Wampler. *Symmetric functions applied to decomposing solution sets of polynomial systems*. Vol. 40, No. 6, pp. 2026-2046. 2002 Society for Industrial and Applied Mathematics.
- [15] A.J. Sommese and J Verschelde. *Numerical homotopies to compute generic points on positive dimensional algebraic sets*. *J. of Complexity* 16(3):572- 602, (2000)
- [16] A.J. Sommese, J Verschelde and C.W. Wampler. (2002a). *A method for tracking singular paths with application to the numerical decomposition* . In algebraic geometry (pp. 329-345). Berlin: de Gruyter.
- [17] A.J. Sommese, J Verschelde and C.W. Wampler. (2001a). *Numerical decomposition of solution sets of polynomial systems into irreducible components*. SIAM J. Number. Anal., 38(6), 2022-2046. NID
- [18] A.J. Sommese, J Verschelde and C.W. Wampler. (2003). *Numerical irreducible decomposition using PHCpack*. In algebra, geometry, and software systems (pp. 109- 129). Berlin: Springer.
- [19] A.J. Sommese, J Verschelde and C.W. Wampler. *Numerical decomposition of the solution sets of polynomial systems into irreducible components*. SIAM J. Numer. Anal. 38(6):2022-2046, 2001.
- [20] A.J. Sommese, J Verschelde and C.W. Wampler. *Numerical irreducible decomposition using projections from points on components*. In *Symbolic Computation: Solving Equations in Algebra, Geometry, and Engineering, volume 286 of contemporary Mathematics*, edited by E.L. Green, S. Hosten, R.C. Laubenbacher, and V. Powers, pages 27- 51. AMS 2001.
- [21] A.J. Sommese, J Verschelde and C.W. Wampler. *Using monodromy to decompose solution sets of polynomial systems into irreducible components*. In *Application of Algebraic Geometry to Coding Theory, Physics and Computation*, edited by C. Ciliberto , F. Hirzebruch, R. Miranda, and M. Teicher. Proceedings of a NATO Conference, February 25- March 1, 2001, Eilat, Israel. Pages 297- 315, Kluwer Academic Publishers.
- [22] A.J. Sommese and C.W. Wampler. *The Numerical Solution of Systems of Polynomials Arising in Engineering and Science*. ISBN 981- 256- 184-6. Word Scientific Publishing Co. Plte. Ltd. 2005 .
- [23] A.J. Sommese and C.W. Wampler. *Numerical algebraic geometry*. In *the mathematics of numerical analysis*. (Park City, UT, 1995), Vol. 32 of lectures in Appl. Math. (pp. 749- 763). Providence, RI: Amer. Math. Soc.
- [24] J. Verschelde(1996). *Homotopy continuation methods for solving polynomial systems*. PhD thesis, Katholieke Universiteit Leuven.
- [25] J. Verschelde(1999). Algorithm 795: PHCpack: A general-purpose solver for polynomial systems by homotopy continuation. ACM Trans. on Math. Software, 25(2), 251- 276.